

Title	Physiologically based pharmacokinetic modelling predicts altered maternal pharmacokinetics of amitriptyline during pregnancy
Authors	Scherf#Clavel, Maike;Leutritz, Anna Linda;Gehrmann, Andrea;Unterecker, Stefan;Walther, Sebastian;Kittel#Schneider, Sarah
Publication date	2025-05-01
Original Citation	Scherf-Clavel, M., Leutritz, A. L., Gehrmann, A., Unterecker, S., Walther, S. and Kittel-Schneider, S. (2025) 'Physiologically based pharmacokinetic modelling predicts altered maternal pharmacokinetics of amitriptyline during pregnancy', British Journal of Clinical Pharmacology. https://doi.org/10.1002/bcp.70084
Type of publication	Article (peer-reviewed);journal-article
Link to publisher's version	10.1002/bcp.70084
Rights	© 2025, The British Pharmacological Society. This is the accepted version of the following item: Scherf-Clavel, M., Leutritz, A. L., Gehrmann, A., Unterecker, S., Walther, S. and Kittel-Schneider, S. (2025) 'Physiologically based pharmacokinetic modelling predicts altered maternal pharmacokinetics of amitriptyline during pregnancy', British Journal of Clinical Pharmacology, which has been published in final form at: https://doi.org/10.1002/bcp.70084 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Download date	2026-09-25 14:38:51
Item downloaded from	https://hdl.handle.net/10468/17556



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Electronic Supplementary Material

to

Physiologically-based pharmacokinetic modeling predicted altered maternal pharmacokinetics of amitriptyline during pregnancy

Short running title: Amitriptyline PBPK model during pregnancy

Maike Scherf-Clavel, PhD¹, Anna Linda Leutritz, MD¹, Andrea Gehrmann, MD¹, Stefan Unterecker, MD^{1,2}, Sebastian Walther, MD¹, Sarah Kittel-Schneider, MD^{1,3,4#}

¹ Department of Psychiatry, Psychosomatics and Psychotherapy, Center of Mental Health, University Hospital of Würzburg, 97080 Würzburg, Germany

² Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, Social Foundation Bamberg, Teaching Hospital of the University of Erlangen, Germany

³ Department of Psychiatry and Neurobehavioural Science, University College Cork, Cork, T12 AK54, Ireland

⁴ APC Microbiome, University College Cork, Cork Ireland

#Correspondence: SKittelSchneider@ucc.ie

Content

1	Clinical Studies.....	3
2	Pregnancy-related enzyme activity changes.....	4
3	Model evaluation	5
3.1	Non-pregnant patients.....	5
3.1.1	Goodness-of-fit plots of predicted vs observed serum concentrations	5
3.1.2	Bias, precision and mean relative of serum concentrations	5
3.1.3	Sensitivity analysis	6
3.2	Pregnant patients	6
3.2.1	Goodness-of-fit plots of predicted vs observed serum concentrations	6
3.2.2	Sensitivity analysis	7
4	Comparison of PK in non-pregnant and pregnant individuals	8
5	References.....	9

1 Clinical Studies

The clinical studies for amitriptyline and nortriptyline model building and evaluation are summarized in table S1.

Table S1 All clinical studies used for amitriptyline model building and evaluation.

Study	PMID	Treatment	Dose	n	n(men/women)	Age (mean±SD) (range)	Dataset	Reference
Amitriptyline iv								
Fridrich et al, 2007	17512256	iv, SD, 1h	25				training	[1]
Fridrich et al, 2007	17512256	iv, SD, 1h	50				training	[1]
Fridrich et al, 2007	17512256	iv, SD, 1h	100				training	[1]
Fridrich et al, 2007	17512256	iv, SD, 2h	25				training	[1]
Schulz et al., 1983	6825390	iv, SD, 2h	43	1	1/0	21	test	[2]
Nortriptyline								
Alvan et al., 1977	852498	im, SD	42	6			training	[3]
Hotha et al., 2010	20853465	po, SD	25	3			test	[4]
Jensen et al., 2012	21999196	po, SD	25	67			test	[5]
Dalen et al., 1998	9585799	po, SD	25	26			test	[6]
Alvan et al., 1977	852498	po, SD	40	6			test	[3]
Amitriptyline po								
Liedholm et al., 1998	9818297	po, SD	25	9	0/9	(25-54)	training	[7]
Mellström et al., 1983	6617075	po, SD	50	9	5/4	(25-49)	training	[8]
Gupta et al., 1998	9597561	po, MD	75	29	19/10	30.2 (19-49)	training	[9]
Venkatakrishnan et al., 2001	11583471	po, SD	50	8	7/1	(24-45)	test	[10]
Gupta et al., 1999	10383563	po, SD	75	15	15/0	29 (21-37)	test	[11]
Nam et al., 2015	25308868	po, SD	10				test	[12]
Nam et al., 2015	25308868	po, SD	25				test	[12]
Balant-Gorgia et al., 1982	7159207	po, SD	75	7	6/1	(22-29)	test	[13]
Baumann et al., 1986	3571944	po, MD	150	30	9/21	45.9	test	[14]
Miljkovic et al., 1996	8980924	po, MD	150	6	3/3	49.8 (31-59)	test	[15]
Miljkovic et al., 1996	8980924	po, MD	75	9	3/6	46.1 (39-56)	test	[15]
Vandel et al., 1993	8436164	po, MD	125	17	7/10	(24-69)	test	[16]
Own_data_50mg (2023-2024)		po, MD	50	269	105/164	55.1±20.0 (18-86)	test	
Own_data_75mg (2023-2024)		po, MD	75	231	89/143	54.6±15.2 (19-81)	test	
Own_data_100mg (2023-2024)		po, MD	100	423	187/236	50.7±17.8 (19-87)	test	
Own_data_125mg (2023-2024)		po, MD	125	86	44/42	48.9±17.0 (18-80)	test	
Own_data_150mg (2023-2024)		po, MD	150	198	113/85	47.0±16.4 (19-79)	test	
Own_data_200mg (2023-2024)		po, MD	200	41	23/18	44.8±12.6 (27-78)	test	

PMID, PubMed-ID; iv, intravenous; SD, single dose; im, intramuscular; po, peroral; MD, multiple dose; n, number; mean±SD, mean±standard deviation

When data on study populations were not available, pharmacokinetics were predicted in a virtual population of 100 individuals. As the study populations were not limited to women of child-bearing age, and pregnant women were not included in the studies, population was build based on a European reference human according to [17]. Mean weight was 68 kg, mean age 53 (predefined age range: 18-100 years) and mean height was 173.10 cm[17]. Mean weight was 68 kg, mean age 53 (predefined age range: 18-100 years) and mean height was 173.10 cm [18].

2 Pregnancy-related enzyme activity changes

Enzyme activity changes due to pregnancy are summarized in table S2.

Table S2 Pregnancy-related enzyme activity changes.

Parameter	Unit	Value used in PBPK model	trimester	Value used in pregnancy model	CYP-activity in pregnancy	Reference
AMITRIPTYLINE						
CYP2C19 Demethylation kcat	1/min	17.37	1. trimester		ND	[19]
			2. trimester	8.685	50% decrease	[19]
			3. trimester	8.685	50% decrease	[19]
CYP2D6 Hydroxylation kcat	1/min	2.71	1. trimester	3.415	ND; 26% increase	[19,21]
			2. trimester	3.66	ND; 35% increase	[19,21]
			3. trimester	4.065	50% increase; 48% increase	[19,21]
NORTRIPTYLINE						
CYP1A2 Demethylation kcat	1/min	0.11	1. trimester	0.0726	33% decrease	[19]
			2. trimester	0.055	50% decrease	[19]
			3. trimester	0.0385	65% decrease	[19]
CYP2C19 Demethylation kcat	1/min	1.55	1. trimester		ND	[19]
			2. trimester	0.775	50% decrease	[19]
			3. trimester	0.775	50% decrease	[19]
CYP2D6 Demethylation kcat	1/min	0.32	1. trimester	0.4032	ND; 26% increase	[19,21]
			2. trimester	0.432	ND; 35% increase	[19,21]
			3. trimester	0.48	50% increase; 48% increase	[19,21]
CYP2D6 Hydroxylation kcat	1/min	4.7	1. trimester	5.922	ND; 26% increase	[19,21]
			2. trimester	6.345	ND; 35% increase	[19,21]
			3. trimester	7.05	50% increase; 48% increase	[19,21]

kcat, catalytic rate constant; ND, No data

3 Model evaluation

3.1 Non-pregnant patients

3.1.1 Goodness-of-fit plots of predicted vs observed serum concentrations




Figure S1 Goodness-of-fit plots for AMI (left) and NOR (right) comparing the predicted serum concentrations with the observed serum concentrations. Line of identity (black solid line), 1.25-fold deviation (black dotted lines), and 2-fold deviation (black dashed lines) are shown.

3.1.2 Bias, precision and mean relative of serum concentrations

Table S3 Bias (mean prediction error), precision (mean absolute prediction error) and mean relative deviation (MRD) of the AMI PBPK model

Dose, Administration	MPE	MAPE	MRD
AMITRIPTYLINE			
10mg, SD	41.14%	56.03%	1.13
25mg, SD	9.82%	36.97%	1.11
50mg, SD	36.94%	60.99%	1.19
75mg, SD	86.07%	101.72%	1.27
75mg, MD	-33.55%	41.65%	1.32
75mg(3x25), MD	-14.30%	14.30%	1.01
125mg, MD	1.40%	14.43%	1.01
150mg, MD	-11.10%	11.90%	1.01
150mg (3x50mg), MD	-1.41%	14.26%	1.01
Mean	12.78%	39.14%	1.12
Min	-33.55%	11.90%	1.01
Max	86.07%	101.72%	1.32
NORTRIPTYLINE			
25mg, SD	-21.97%	51.51%	1.35
50mg, SD	-1.70%	42.75%	1.39
75mg, SD	41.42%	66.67%	1.26
75mg, MD	10.23%	14.40%	1.01
75mg(3x25), MD	-12.23%	12.23%	1.01
125mg, MD	6.61%	6.61%	1.00
150mg, MD	22.98%	22.98%	1.02
150mg (3x50mg), MD	4.07%	13.64%	1.01
Mean	6.18%	28.85%	1.13
Min	-21.97%	6.61%	1.00
Max	41.42%	66.67%	1.39

MPE; mean prediction error; MAPE, mean absolute prediction error; MRD, mean relative deviation; SD, single dose; MD, multiple dose; Min, minimum; Max, maximum

3.1.3 Sensitivity analysis




Figure S2 Sensitivity for parameters that were estimated during model building was measured as relative changes in c_{trough} of multiple dose administration (75mg) of AMI. Variation range was range was 10.0 with maximum number of steps = 9.

3.2 Pregnant patients

3.2.1 Goodness-of-fit plots of predicted vs observed serum concentrations




Figure S3 Goodness-of-fit plots for AMI (left) and NOR (right) comparing the predicted serum concentrations with the observed serum concentrations in pregnancy. Line of identity (black solid line), 1.25-fold deviation (black dotted lines), and 2-fold deviation (black dashed lines) are shown.

3.2.2 Sensitivity analysis




Figure S4 Sensitivity for parameters that were estimated during model building, as well as the 9 pregnant specific compartments and the new included CYP2D6 hydroxylation of amitriptyline was measured as relative changes in c_{trough} of multiple dose administration (75mg) of AMI during pregnancy. Variation range was range was 10.0 with maximum number of steps = 9.

4 Comparison of PK in non-pregnant and pregnant individuals

Comparison of median and IQR of trough serum concentrations for amitriptyline, nortriptyline and the active moiety in non-pregnant, as well as pregnant women are shown in figure S5.




Figure S5 Comparison of median and IQR of trough serum concentrations for amitriptyline, nortriptyline and the active moiety in non-pregnant, as well as pregnant women.

5 References

1. Fridrich P, Colvin HP, Zizza A, Wasan AD, Lukanich J, Lirk P, Saria A, Zernig G, Hamp T, Gerner P. Phase 1A safety assessment of intravenous amitriptyline. *J Pain* 2007; 8: 549-555
2. Schulz P, Turner-Tamiyasu K, Smith G, Giacomini KM, Blaschke TF. Amitriptyline disposition in young and elderly normal men. *Clin Pharmacol Ther* 1983; 33: 360-366
3. Alván G, Borga O, Lind M, Palmér L, Siwers B. First pass hydroxylation of nortriptyline: concentrations of parent drug and major metabolites in plasma. *Eur J Clin Pharmacol* 1977; 11: 219-224
4. Hotha KK, Ravindranath LK, Veera KN, Kumar KK, Reddy YR, Jagadeesh B, Rao DV, Bharathi DV, Mullangi R. Development and validation of a highly sensitive LC-MS/MS method for simultaneous quantitation of nortriptyline and 10-hydroxynortriptyline in human plasma: application to a human pharmacokinetic study. *Biomed Chromatogr* 2010; 24: 1113-1119
5. Jensen BP, Roberts RL, Vyas R, Bonke G, Jardine DL, Begg EJ. Influence of ABCB1 (P-glycoprotein) haplotypes on nortriptyline pharmacokinetics and nortriptyline-induced postural hypotension in healthy volunteers. *Br J Clin Pharmacol* 2012; 73: 619-628
6. Dalén P, Dahl ML, Bernal Ruiz ML, Nordin J, Bertilsson L. 10-Hydroxylation of nortriptyline in white persons with 0, 1, 2, 3, and 13 functional CYP2D6 genes. *Clin Pharmacol Ther* 1998; 63: 444-452
7. Liedholm H, Lidén A. Food intake and the presystemic metabolism of single doses of amitriptyline and nortriptyline. *Fundam Clin Pharmacol* 1998; 12: 636-642
8. Mellström B, Bertilsson L, Lou YC, Säwe J, Sjöqvist F. Amitriptyline metabolism: relationship to polymorphic debrisoquine hydroxylation. *Clin Pharmacol Ther* 1983; 34: 516-520
9. Gupta SK, Shah J, Guinta D, Hwang S. Multiple-dose pharmacokinetics and pharmacodynamics of OROS and immediate-release amitriptyline hydrochloride formulations. *J Clin Pharmacol* 1998; 38: 60-67
10. Venkatakrishnan K, Schmider J, Harmatz JS, Ehrenberg BL, von Moltke LL, Graf JA, Mertzanis P, Corbett KE, Rodriguez MC, Shader RI, Greenblatt DJ. Relative contribution of CYP3A to amitriptyline clearance in humans: in vitro and in vivo studies. *J Clin Pharmacol* 2001; 41: 1043-1054
11. Gupta SK, Shah JC, Hwang SS. Pharmacokinetic and pharmacodynamic characterization of OROS and immediate-release amitriptyline. *Br J Clin Pharmacol* 1999; 48: 71-78
12. Nam Y, Lim CH, Lee HS, Chung SJ, Chung YH, Shin YK, Kim MG, Sohn UD, Kim HC, Jeong JH. Single-dose, randomized, open-label, 2-way crossover study of the pharmacokinetics of amitriptyline hydrochloride 10- and 25-mg tablet in healthy male Korean volunteers. *Clin Ther* 2015; 37: 302-310
13. Balant-Gorgia AE, Schulz P, Dayer P, Balant L, Kubli A, Gertsch C, Garrone G. Role of oxidation polymorphism on blood and urine concentrations of amitriptyline and its metabolites in man. *Arch Psychiatr Nervenkr (1970)* 1982; 232: 215-222
14. Baumann P, Jonzier-Perey M, Koeb L, Lê PK, Tinguely D, Schöpf J. Amitriptyline pharmacokinetics and clinical response: I. Free and total plasma amitriptyline and nortriptyline. *Int Clin Psychopharmacol* 1986; 1: 89-101
15. Miljković B, Pokrajac M, Timotijević I, Varagić V. Clinical response and plasma concentrations of amitriptyline and its metabolite-nortriptyline in depressive patients. *Eur J Drug Metab Pharmacokinet* 1996; 21: 251-255
16. Vandel S, Bertschy G, Perault MC, Sandoz M, Bouquet S, Chakroun R, Guibert S, Vandel B. Minor and clinically non-significant interaction between toloxatone and amitriptyline. *Eur J Clin Pharmacol* 1993; 44: 97-99
17. [Anonymous]. Basic anatomical and physiological data for use in radiological protection: reference values. A report of age- and gender-related differences in the anatomical and physiological characteristics of reference individuals. ICRP Publication 89. *Ann ICRP* 2002; 32: 5-265

18. Open System Pharmacology. PK-Sim®. Version 11.0. Available online:.
19. Anderson GD. Pregnancy-induced changes in pharmacokinetics: a mechanistic-based approach. *Clin Pharmacokinet* 2005; 44: 989-1008
20. Tylutki Z, Mendyk A, Polak S. Mechanistic Physiologically Based Pharmacokinetic (PBPK) Model of the Heart Accounting for Inter-Individual Variability: Development and Performance Verification. *J Pharm Sci* 2018; 107: 1167-1177
21. Abduljalil K, Furness P, Johnson TN, Rostami-Hodjegan A, Soltani H. Anatomical, physiological and metabolic changes with gestational age during normal pregnancy: a database for parameters required in physiologically based pharmacokinetic modelling. *Clin Pharmacokinet* 2012; 51: 365-396